

## Cultivated eggplants – origin, breeding objectives and genetic resources, a review

*Agnieszka Sękara, Stanisław Cebula, Edward Kunicki*

Department of Vegetable Crops  
Agricultural University in Kraków  
29 Listopada 54, 31-425 Kraków, Poland  
e-mail: asekara@ogr.ar.krakow.pl

Key words: *Solanum melongena*, *S. aethiopicum*, *S. macrocarpon*, taxonomy, cultivars, genetic diversity

### ABSTRACT

Brinjal eggplant (*Solanum melongena* L.) is an agronomically important non-tuberous crop of *Solanaceae* family, native to southern India, and widely grown in Americas, Europe and Asia. Its close relatives: *S. aethiopicum* (scarlet eggplant) and *S. macrocarpon* (Gboma eggplant) are of African origin. Brinjal eggplant is a new vegetable in Poland. Until recently it was grown only in glasshouse conditions. Breeding of new eggplant cultivars and improvement of growing technology resulted in the increase of both cultivation acreage and consumption scale in Poland. For an efficient development of eggplant breeding programmes it is necessary to collect information concerning the characteristics and variety of eggplant genetic resources. The present publication includes detailed taxonomy of eggplant and its relatives, their origin and history of growing. The information connected with eggplant worldwide genetic resources and breeding objectives have been presented. Different propositions for cultivar grouping and the characteristics of selected cultivars have been taken into consideration.

### Taxonomy

*Solanaceae* is a plant family comprising about 2300 species, nearly one-half of which belong to the genus *Solanum*. Most species within this genus are endemic to the Americas, only 20% are Old World species. Family *Solanaceae* has been the source of many morphologically different domesticated species. Doganlar et al. (2002b) explained this phenomenon with the fact that only a few conserved major loci are responsible for the dramatic phenotypic changes that accompanied domestication in this family. Like tomato and pepper, eggplant is an autogamous diploid with 12 chromosomes. A molecular genetic linkage map based on tomato cDNA, genomic DNA, and EST markers was constructed for *S. melongena* (Doganlar et al. 2002a, SOL Genomic Network 2006). The comparison of the eggplant and tomato maps revealed that chromosomal evolution since their divergence from the common ancestor has proceeded primarily through inversion of otherwise conlinear segments of the genome. Overall, eggplant and tomato are differentiated by 28 rearrangements, which could be explained by 23 paracentric inversions and five translocations during evolution from the last common ancestor of the species. Eggplant and tomato are diverged three- to six fold more than tomato and potato (Fig. 1). The domestication of the *Solanaceae* has been driven by mutations in a very limited number of target loci with major phenotypic effect. Selection pressures were exerted on the same loci despite independent domestication of the crops on different continents. The conservation of gene function among the *Solanaceae* indicates that the wealth of knowledge of genetic analysis of tomato, potato and pepper can be used as a springboard for rapid advancement of eggplant genetics (Frery et al. 2003).




Fig. 1. Dendrogram of crop plants from *Solanaceae* family (Olmstead and Palmer 1997)

The genomics in genus *Solanum* are evolving at a moderate pace compared to other plant species (Doganlar et al. 2002a). The species of *Solanum* subgenus *Leptostemonum* comprise almost one third of the genus and are distributed worldwide. The members of this monophyletic subgenus are defined by their sharp epidermal prickles, thus they are commonly referred to as the ‘spiny solanums’. The subgenus *Leptostemonum* includes a number of economically important species such as the cultivated eggplants (Levin et al. 2005, Levin et al. 2006).

The common name ‘eggplant’ encompasses three closely related cultivated species, endemic to the Old World, belonging to the genus *Solanum* L., subgenus *Leptostemonum* (Dunal) Bitter and two sections:

1. Section *Melongena*

- *S. melongena* L. (brinjal eggplant, aubergine) – synonyms: *Solanum cumingii* Dunal, *Solanum pressum* Dunal, *Solanum undatum* Poiret sensu Ochse;
- *S. macrocarpon* (Gboma eggplant) – synonyms: *Solanum integrifolium* Poiret var. *macrocarpum*, *Solanum melongena* L. var. *depressum* Bail.

2. Section *Oliganthes*

- *S. aethiopicum* L. (scarlet eggplant, jaxatu) – synonyms: *Solanum integrifolium* Poiret, *Solanum integrifolium* Poiret var. *microcarpum*, *Solanum gilo* Raddi, *Solanum naumannii* Engl., *Solanum pierreanum* Pailleux & Bois, *Solanum zuccagnianum* Dunal.

Historical and modern attempts of eggplant and its relatives’ classification were in detail presented by Mace et al. (1999). The historical difficulties and confusions connected with *Solanum* species classification is reflected by the fact that 1000-1400 *Solanum* species have been described under more than 3000 binominal names (Daunay and Lester 1988). Proper classification of eggplant cultivars and accessions collected all over the world is possible to achieve with the use of molecular methods, such as supplementation of morphological analyses. The other parameters (morphological features, crossability, F<sub>1</sub> fertility) are insufficient, because of the high morphological variability in large and complex genus *Solanum*, and the possibility of crossing between distantly related species (Daunay and Lester 1988, Furini and Wunder 2003).

According to Karihaloo et al. (2002), all members of the eggplant complex were revealed to bear very high similarity to each other, based on studies of the seed protein profiles by SDS-PAGE. On the other hand, Furini and Wunder (2003) reported that the eggplant complex showed a very large morphological variation, to some extent reflected in the unrooted tree based on the AFLP data. The authors explained such a high degree of variation by using, in their analyses, the accessions derived from Asian countries where the greatest diversity is found. Also, their DNA analyses were based on AFLP markers which proved to be more informative than RAPD markers and allozymes. Furini and Wunder (2003) suggested that

while all members of the eggplant aggregate cluster together, sufficient genomic flexibility has been created within the group to adapt to changes in the environment.

*S. melongena* may have been indirectly derived from the wild *S. incanum*, domesticated in India and Southeast of China. *S. aethiopicum* and *S. macrocarpon* were domesticated in Africa from their wild relatives *S. anguivi* and *S. dasyphyllum* (Lester 1998).




Fig. 2. Dendrogram of *Solanum* species by UPGMA based on the genetic distance of Nei and Li (Sakata and Lester 1997)

From the dendrogram constructed by the unweighted pair-group method, Sakata and Lester (1997) suggested that *S. incanum* is a species close to *S. melongena*, and the next close species is *S. macrocarpon* (Fig. 2). High genetic identities between *S. melongena*, *S. incanum* (wild form) and *S. insanum* (weedy form) were proved by Karihaloo and Gottlieb (1995) on the base of enzyme electrophoretic studies, and Karihaloo et al. (1995) on the base of the random amplified polymorphic DNA (RAPD) variation. Close genetic relationship between *S. melongena* and *S. incanum* was confirmed by Singh et al. (2006) using RAPD technique. *S. melongena* and *S. macrocarpon* were included to the section *Melongena* and *S. aethiopicum* to the section *Oliganthes* (Doganlar et al. 2002a). The results of Karihaloo et al. (2002) based on the seed protein study, and Furini and Wunder (2003) based on the amplified fragment length polymorphism (AFLP) markers indicate that *S. macrocarpon* is more related to *S. aethiopicum* (section *Oliganthes*) than to *S. melongena*. Similar results were obtained by Mace et al. (1999) based on the AFLP analysis of genetic relationship among the cultivated eggplants and wild relatives. The authors confirmed the close relationship between *S. macrocarpon* and *S. dasyphyllum*. The fact that *S. macrocarpon* and *S. aethiopicum* were domesticated and are cultivated mainly in Africa supports their relatively similar topology in the dendrogram (Furini and Wunder 2003).

*Origin, domestication and economic importance*

*S. melongena* occurs in wild or semi wild form in India. Various data indicate, that from the several species that evolved in Africa, one (*S. incanum*), gave rise to a distinct species which spread to South-East Asia as the wild ancestor of *S. melongena* (Lester 1998). India or Indochina are recognized as the centre of the eggplant diversity. Primitive eggplant characters are tall plants with large, spiny leaves, flowering in clusters with andromonoecy. Their fruits are small, green, and bitter in taste, with thick skin and hard flesh. Domestication, mutation, natural intercrossing, human selection and hybridization brought extensive genetic diversity of eggplant cultivars grown all over the world. Cultivar differences concern mainly the colour, shape and height of fruits, but chemical composition of fruits, the earliness of fruiting, yielding, environmental requirements, etc. are also taken into consideration. Fruit colour varies from light to dark purple, almost black, green, or white. Fruit length is between 4-45 cm, and thickness 2-35 cm, at different shapes and weight ranging between 15-1500 g. The fruits are set as single or in clusters, up to 5 fruits. Physiologically ripe fruits become brown, red or yellow (Swarup 1995).

*S. aethiopicum* is a fruit and leaf vegetable. It is a herbaceous shrub with hairy or glabrous leaves and hermaphroditic flowers, self or cross pollinated, single or in clusters. The fruits are consumed raw or cooked. They are light to dark green, white or blackish in colour, with a bitter taste that varies depending on saponin content. The fruits shape is round, elongate-round or oval with smooth or grooved surface and taste varying from sweet to bitter, particularly in the case of oval-fruit cultivars. At full maturity, the fruits turn red or reddish-orange due to high carotene content. Fruit surfaces vary from smooth to grooved or ribbed. The leaves are often consumed in the same way as spinach (Seek 1997, Macha 2005).

*S. macrocarpon* is grown for large, glabrous leaves (50 × 30 cm), used as a green vegetable. Fruits have large, often clasping calyx. They are sub spherical and large (3-10 cm in diameter, 2-6 cm long), cream white, green-white or green. Fruits are sweeter in taste (in comparison to *S. aethiopicum*), and most preferred. At full maturity fruits turn yellow, orange or brown with cracked surface (Bukanya 1994, Macha 2005).

Brinjal eggplant was described in India in 3<sup>rd</sup> century B.C. There were at least 33 sanskrit names for eggplant in ancient Indian literature, the most common being Varttaka, Bhandaki and Nattingan (Swarup 1995). The growing of eggplant small-fruited characters began in 4<sup>th</sup> century in China, and in 9<sup>th</sup> century in Africa. Although cultivated from prehistorical times, eggplant appears to have been unknown to the Western World for many centuries. Numerous Arabic and African names for eggplant, and the lack of ancient Greek and Roman names indicate that the vegetable was carried into the Mediterranean by the Arabs relatively late, probably in 7<sup>th</sup> century. *Melongena* was an Arabic name for one of eggplant

cultivars. Avicenna, “the father of modern medicine”, mentioned it as a medicinal and vegetable plant. Eggplant was established in Poland by 15<sup>th</sup> century, initially as ornamental and medicinal species.

At present brinjal eggplant is the third, after potato and tomato, most important crop from *Solanaceae* family. Greatest eggplant producers are China (17 mln tons per year), India (8 mln tons), Egypt (1 mln tons) and Turkey (0.9 mln ton) (FAOSTAT Data 2006). In Poland, like in many Central European countries, eggplant is still an exotic vegetable but in Asia and the Mediterranean it is an important and valuable nourishment component, the so-called ‘the king of vegetables’.

African eggplants – *S. aethiopicum* and *S. macrocarpon*, are the most popular native, traditional vegetables in West and Central Africa, but the productivity of these crops is still relatively low and the growing area and yields have not been statistically evaluated. The centre of these eggplants diversity is Western Africa. African eggplants are grown mainly in gardens and small fields near villages. *S. aethiopicum* is widely grown in South America, and *S. macrocarpon* in Asia and tropical America, too.

#### *Genetic resources*

Genetic resources of eggplants have been collected methodically only in some European and Asian countries, and there is no comprehensive collection dedicated to the germplasm of eggplants available worldwide. Gene banks for eggplant and related species have been collected by the partners of EGGNET project in Europe: INRA-Montfavet, France; Botanical and Experimental Garden, University of Nijmegen, The Netherlands; Birmingham University Botanic Gardens, Birmingham, United Kingdom; Plant Research International (PRI), Centre for Genetic Resources (CGR), Wageningen, The Netherlands; Biotechnology Department (Genetics), Polytechnic University of Valencia (UPV) Valencia, Spain; Department of Vegetables, Agricultural Research Centre of Macedonia and Thrace (ARCMT), Themi-Thessaloniki, Greece; Istituto del Germoplasma (IDG), Consiglio Nazionale delle Ricerche, Bari, Italy; Institute of Plant Genetic and Crop Plant Research (IPK), Gatersleben, Germany, and Institut National d’Horticulture (INH), Angers, France. EGGNET connects all the European collections and creates one European collection, including more than 1700 accessions of *S. melongena* L., 750 accessions of *S. aethiopicum* and 210 accessions of *S. macrocarpon* (Daunay et al. 2003, 2006, ECP/GR Eggplant Database 2006).

The greatest gene banks of eggplants in other world regions have been collected, among the others, in the National Germplasm Resources Laboratory in Beltsville, Maryland, USDA / ARS National Genetic Resources Program (GRIN) – 770 accessions of *S. melongena*, 60 of *S. aethiopicum* and 4 of *S. macrocarpon*; National Bureau of Plant Genetic Resources, New Delhi, India; Chinese Academy

of Agricultural Sciences, Institute of Crop Germplasm Resources, National Gene Bank of China (NGBC) – 1300 accessions of *S. melongena*; National Institute of Agrobiological Sciences, Japan – 561 accessions of *S. melongena* and 31 of *S. aethiopicum*; Asian Vegetable Research and Development Center, Shanhua, Taiwan – 1777 accessions of *S. melongena*, 30 of *S. aethiopicum* and 133 of *S. macrocarpon*, and Vavilov Research Institute of Plant Industry, St. Petersburg, Russia – 238 accessions of *S. melongena* (Mueller et al. 2005, SOL Genomic Network 2006).

### *Breeding objectives*

*S. melongena* is an object of intensive breeding programs mainly in the countries with intensive production of this species (Western Europe, Turkey, India, China, Japan). Many F<sub>1</sub> hybrids, with differentiated phenotypes are the result of breeding works held in the last thirty years. New *S. aethiopicum* cultivars are bred in some African countries, but on a small scale, due to economic, sociological, and political situation in this region. *S. macrocarpon* cultivars are improved locally by growers, mainly in Africa.

Cultivated *S. melongena* genotypes often have insufficient levels of resistance to biotic and abiotic stresses. Genetic resources of this species have been assessed for resistance to its most serious diseases and pests. The attempts at crossing eggplant with its wild relatives resulted in limited success due to sexual incompatibilities. Eggplant tissues present a high morphogenetic potential, useful for developmental studies as well as establishing biotechnological approaches to produce improved varieties, such as embryo rescue, *in vitro* selection, somatic hybridization and genetic transformation (Collonnier et al. 2001, Magioli and Mansur 2005). Eggplant *in vitro* regeneration can be induced from different explants with the use of distinct growth regulators and morphogenetic pathways, as it was summarized by Magioli and Mansur (2005). For induction of embryogenesis or organogenesis hypocotyl, epicotyl, cotyledon, stem nodes, leaf, and roots were used (Sharma and Rajam 1995, Magioli et al. 1998, Magioli et al. 2001, Franklin and Lakshmi Sita 2003). In addition, the availability of efficient transformation protocols favors gene regulation studies, in particular those related to embryogenesis, with advantages over other species. From this perspective, eggplant can be considered as alternative model plant for studying different aspects of plant biology (Magioli and Mansur 2005). Kashyap et al. (2003) summarized the efforts to improve eggplant genetics with emphasis on the use of biotechnology to introgression genes from wild species into cultivated eggplant. For example, populations derived from the cross between *S. linnaeanum* and *S. melongena* may be useful for future mapping of disease resistance and abiotic stress loci and for transfer of these resistances to cultivated eggplant (Frary et al. 2003). Tolerance to

*Verticillium* wilt was found in a wild relative of eggplant, *Solanum torvum*. To transfer this tolerance to eggplant (*S. melongena*), protoplast fusions between eggplant and irradiated *S. torvum* protoplasts were performed by Jarl et al. (1999).

The main *S. melongena* breeding objectives are:

- resistance to diseases, insects and nematodes (Fusarium wilt *Fusarium oxysporum*, Verticilium wilt *Verticillium dahliae*, Bacterial wilt *Pseudomonas solanacearum*, eggplant fruit borer *Leucinodes orbonalis*, melon aphid *Aphis gossypii*, leafhopper *Amrasca biguttula*, root-knot nematode *Meloidogyne* spp., etc.),
- resistance or tolerance to abiotic stress (drought, low or high temperatures, salinity),
- parthenocarpy,
- resistance to herbicides,
- yield and its quality (the world average 15.7 t ha<sup>-1</sup>, Africa 19.2 t ha<sup>-1</sup>, North America 23.1 t ha<sup>-1</sup>, South America 13.1 t ha<sup>-1</sup>, Asia 15.1 t ha<sup>-1</sup>, Europe 28.3 t ha<sup>-1</sup>. The yield of F<sub>1</sub> hybrids often exceeds 40-50 t ha<sup>-1</sup> and is characterized by early maturity, uniform harvest, rich colour, taste and aroma, and great storage quality),
- nutritive value (high dry matter, sugars, anthocyanin and total phenol contents, low level of polyphenol oxidase activity and orthodihydroxy phenolic compounds to avoid browning of cut fruits),
- market needs and consumer preference (Swarup 1995, Farooqui et al. 1997, Kumar et al. 1998, Donzella et al. 2000, Acciarri et al. 2002, Prabhavathi et al. 2002, Rahman et al. 2002, Pessarakli and Dris 2004, Sidhu et al. 2005).

Some specific objectives of *S. aethiopicum* breeding programs are: developing of hybrids and open pollinated cultivars resistant to mites, *Stemphylium* and *Alternaria* complex; improvement of yield and fruit quality (colour, size, shape etc.) (Anaso et al. 1990, Seek 1997). Omidiji (1983) evaluated the cultivars and F<sub>1</sub> hybrids of *S. aethiopicum* for the yield and its components in Nigeria. The results of breeding for high yield of berries and improved yield attributes of homozygous lines of *S. aethiopicum* were presented by Anaso et al. (1990). Carvalho and Ribeiro (2002) presented the results of combining ability analysis in diallel crosses among three Brazilian *S. aethiopicum* cultivars.

*S. aethiopicum* and *S. macrocarpon* are of interest for genetic improvement of *S. melongena*, considering the presence of several traits of agronomic interest in its germplasm (Alba et al. 2005). Daunay et al. (1993) reported the possibility of production and characterization of fertile somatic hybrids of *S. melongena* and *S. aethiopicum* Aculeatum group. Schaff et al. (1982) suggested the possibility of the transfer of genes for resistance to two-spotted spider mite from *S. macrocarpon* to *S. melongena*. The interspecific hybridisation was used to transfer the resistance to *Pseudomonas solanacearum* from *Solanum aethiopicum* to *S. melongena*. With this



method it was possible to obtain a large variation in the shape and colour of the fruits (Ano 1989). Gowda et al. (1990) used the interspecific hybridisation between *S. macrocarpon* and *S. melongena* to obtain hybrids resistant to shoot and fruit borer (*Leucinodes orbonalis*). Rizza et al. (2002) obtained dihaploid plants through anther culture of somatic hybrids between *S. melongena* and *S. aethiopicum* Gilo group which were characterized by complete resistance to fungal wilt caused by *Fusarium oxysporum* f. sp. *melongenae*.

#### Cultivars

*S. melongena* is characterized by great morphological diversity. Martin and Rhodes (1979) cited the classification proposed by Choudhury (1976) who divided brinjal eggplant cultivars into three botanical varieties on the basis of the shape of the fruit:

- *S. melongena* var. *esculentum* Dunal (Nees) – round, oval or egg-shaped fruits,
- *S. melongena* var. *serpentinum* L. – long slender fruits,
- *S. melongena* var. *depressum* L. – small, miniature fruits, dwarf, and early types.

Martin and Rhodes (1979) classified 475 eggplant cultivars from the world collection assembled by the U.S. Department of Agriculture into 11 groups, by 18 characteristics, using numerical taxonomic methods:

- I. Tubefrut – dark purple, long fruits, region of origin from India to Turkey,
- II. Spinigrene – spiny foliage, greenish fruits – from India and Pakistan,
- III. Cluster – fruits in clusters, often white and long,
- IV. Oldster – long-lived plants with greenish or light purple fruits – from India and Pakistan,
- V. Pretiblum – dark, purple flowers – from India and Pakistan,
- VI. Talgro – low yields, long, dark purple fruits – from Turkey,
- VII. Stipe – streaked fruits – from India to Turkey,
- VIII. Earlymot – high yields, early flowering – from India to Turkey,
- IX. Lakluster – early flowering, low vigor, mottled fruits – from India to Turkey,
- X. Lateg – fruits often ovoid, late flowering – from India to Turkey,
- XI. Violeaf – dark purple foliage and stems, long, dark purple fruits – unclear origin.

The cited authors found that some similarity groups were composed of cultivars from various countries from India to Turkey. This fact suggests a wide distribution of the principal character associations. Also cultivars from Europe, Africa and New World tended to fall into several different similarity groups, which suggests their diverse origins. Geographical groups can be characterized, in some cases, by distinctive characteristics as determined from group means, which clearly reflect preferences in the case of some countries, i.e. USA or Japan (Martin and Rhodes 1979).

Prohens et al. (2005) studied the molecular and morphological diversity of 28 Spanish traditional cultivars of eggplant, collected in four groups: “round”, “semi-long”, “long”, and “listada de Grandia”. The authors found that eggplant cultivars groups showed some genetic differences, and no individual AFLP (amplified fragment length polymorphism) markers specific and universal to one cultivar group could be found. “Round” cultivars were genetically more diverse than other groups. The results suggested that evolution of eggplants in Spain has involved frequent hybridizations and frequent movement and exchange seeds. The structure and diversity among the regions indicate that most of the diversity can be collected in single selected regions (Prohens et al. 2005).

The mentioned types of eggplant cultivars, differentiated on the base of the shape and colour of the fruit and the origin, are often used in popular and professional literature.

Dark purple eggplants (western eggplants) are typical for America and Europe, but introduced into Asia. Plants are less vigorous but very productive. Fruits come in two basic shapes, oval and elongated and are usually large (200-600 g).

- oval: ‘Black Beauty’, ‘Bonica F<sub>1</sub>’, ‘Classic F<sub>1</sub>’, ‘Epic F<sub>1</sub>’, ‘Galine F<sub>1</sub>’, ‘Sonata F<sub>1</sub>’, ‘Nadia F<sub>1</sub>’, ‘Tudela F<sub>1</sub>’, ‘Gostbuster F<sub>1</sub>’(white), ‘Listada de Grandia’ (purpure striped with white), ‘Neon F<sub>1</sub>’ (pink purple), ‘Rosita’ (pink lavender), ‘Zebra F<sub>1</sub>’ (violet striped with white).
- elongated: ‘Baluroi F<sub>1</sub>’, ‘Black King’, ‘Fabina F<sub>1</sub>’, ‘Ichiban F<sub>1</sub>’, ‘Long Purple’, ‘New Purple’, ‘Nite Lady F<sub>1</sub>’, ‘Lavender Tough F<sub>1</sub>’ (white), ‘Casper’ (white), ‘Cloud Nine F<sub>1</sub>’ (white), ‘Antigua’ (white striped with lavender), ‘Fairy Tale F<sub>1</sub>’ (lavender with white strikes), ‘Louisiana Long Green’ (green).

Miniature eggplants (Italian, baby, finger eggplants) – few inches long, narrow or rounded; they are generally sweeter and more tender than the larger varieties; they also have thinner skins and contain fewer seeds (‘Baby Bell’ – black-purple, ‘Bianca Rosa’ – white with lavender streaks, ‘Prosperosa’ – cherry eggplant, round, violet, ‘Violetta di Firenze’ – violet).

Oriental eggplants, native to tropical Asia, are very popular in Japan, China, India, Thailand and the Philippines. Plants are early and vigorous. Fruits could be purple, violet green, with or without stripes; round or slender in shape. Tender and sweet in taste, they are cooked without peeling or salting, stuffed or baked in India, tempura in Japan, stir-fried or boiled in China. In South-eastern Asia many primitive varieties are cultivated, with small green fruits and slight anthocyanin pigmentation. Oriental eggplants are divided into the following groups (SeedQuest 2006):

- Chinese eggplants – usually long, violet, purple lavender-bluish white, especially low in seeds, tender and sweet, plants are vigorous and prolific (‘Little Fingers F<sub>1</sub>’ – purple, ‘Long White Angel F<sub>1</sub>’ – white, ‘Ma Zu Purple’ – purple, ‘Megadok F<sub>1</sub>’ – purple, ‘Ping Tung Long’ – dark-purple, ‘Lucky Green’ – green,

- Japanese eggplants are firmer and heavier, sweet in taste, violet to inky-purple, long or egg-shaped ('Kurume Long Purple' – dark purple, 'Kyoto Egg F<sub>1</sub>' – dark purple, round, 'Milionaire F<sub>1</sub>' – dark purple, long, 'Senryu Ni Gou F<sub>1</sub>' – dark purple, elongated, 'Shoya Long F<sub>1</sub>' – dark purple, very long,
- Thai eggplants come in two groups: small, round, tomato-like ones (40-80 g – 'Green Doll F<sub>1</sub>' – white with green stripes, 'Kermit F<sub>1</sub>' – green/white, 'Thai Round Green' – green, 'Violet Prince F<sub>1</sub>' – violet, 'White Ball F<sub>1</sub>' – white, 'Puangyok Thai Pea' – green cherry eggplant) or very elongated (10 inches, about 100-200 g – 'Rolex F<sub>1</sub>' – green, 'Tai Long Green F<sub>1</sub>' – green). They are mild and sweet, dense, with thick skin, and many seeds. Plants bearing lots of fruits are very attractive and often grown as ornamentals,
- Indian eggplants – small 30-100 g, oval, violet to purple, sometimes with stripes ('Apsara F<sub>1</sub>' – purple with white stripes, 'Chu Chu F<sub>1</sub>' – red purple, 'Manjari', 'Ratna' – dark purple, 'Rhim Jhim' – purple with white streaks, 'Hari F<sub>1</sub>' – long, green, 'Bali F<sub>1</sub>' – long, purple) or medium sized (200-300 g) ('Bharata Star F<sub>1</sub>' – round, dark purple, 'Suphal' – dark purple, oval, 'Supriya' – round, violet).

Only two eggplant cultivars have been entered into The Polish National List of Vegetables: 'Classic F<sub>1</sub>' (Clause Tézier) and 'Epic F<sub>1</sub>' (Seminis Vegetable Seeds, Inc.) and one cultivar applied for addition to the NLI and accepted for testing '2RGD-86' (IW, Skierniewice) (COBORU, 2006).

*S. aethiopicum* is a phenotypically diverse species. *S. aethiopicum* cultivars could be divided into four groups:

- Gilo Group – with hairy, inedible leaves, and fruits differentiated in shape (round, elongated, egg-shaped or spindly), colour and size (from a few to several dozen grams),
- Shum Group – with glabrous leaves eaten as a green vegetable and very small, inedible, small, elongated fruits,
- Kumba Group – with glabrous, large leaves, eaten as a green vegetable, 5-10 cm in diameter fruits also edible,
- Aculeatum Group – ornamentals with hairy leaves and large fruits, often used for disease resistance breeding under the synonym *S. integrifolium* (Lester 1986, Caruso 2001, Lester and Daunay 2003).

The Gilo cultivar group might have evolved from the Shum cultivar group through hybridization and selection (Anaso 1991). Stedje and Bukenya-Ziraba (2003) studied the variation in 39 polymorphic RAPD markers for 18 populations of *S. anguivi* and *S. aethiopicum* Gilo and Shum Group. The variation among the species and groups was less than 10%, whereas the variation within the species and groups was more than 90%. The Gilo and Shum Groups of *S. aethiopicum* kept morphologically distinct by a strong man-made selection pressure.

Superior varieties of *S. aethiopicum* have yet to be identified or developed. The researches on new, promising varieties are led at the African branch of the Asian Vegetable Research and Development Center (AVRDC). The best known indigenous cultivars are: 'Manyire Green' – a popular cultivar of East Africa, characterized with round, green ripening to red fruits, 'Tengeru White' – round, half green, half white fruits, 'Jaxatu Soxna' – flat, ribbed fruits, light green to white, 50-80 g, 'N'Goyo' – flat, ribbed, dark-green fruits, 70-80 g, 'N'Galam' – flat, ribbed, light-green fruits, 120-180 g. In Europe and Americas the most widely grown are: 'Turkish Orange' – small, orange spherical fruits, about 6 cm in diameter, with an excellent sweet taste, 'Sweet Red' – small, attractive fruits, about 2.5 cm in diameter, with the green skin with dark-green stripes, turning red with dark-red stripes at maturity, 'Small Ruffled Red' – small, deeply creased fruits, about 5 cm in diameter, with orange-red skin, 'Comprido Verde Claro' and 'Morro Redondo' – small ribbed fruits in a bright, acid green, widely grown in Brazil, 'Sweet African Egg' with small, gently ribbed white fruits that mature to a glowing orange, both decorative and tasty.

*S. macrocarpon* is less morphologically diverse than *S. melongena* and *S. aethiopicum*. Bukenya and Carasco (1994) recognized four groups of *S. macrocarpon* complex in Uganda: *S. macrocarpon* (semi wild group), *S. macrocarpon* 'Mukono' cv. Group and *S. macrocarpon* 'Nabingo' cv. Group. With the use of morphological and experimental methods they found the considerable variation within the two major cultivar groups, in particular 'Munoko'. The variation within *S. macrocarpon* complex is attributable to genotypic differences and environmental factors (Bukenya and Hall 1987).

Eggplants are species of economic importance mainly in Asian and African countries. They are widely cultivated in America and Europe. Professional growers see the necessity of introducing new varieties, particularly hybrids, breeding for very high quality production, in different climatic zones. Eggplants are used for the treatment of many diseases i.e. diabetes, bronchitis, asthma, dysuria, dysentery, etc. For an effective breeding program, information concerning the extent and nature of genetic diversity within the crop species is essential (Singh al. 2006). It is useful for characterizing individual accessions and cultivars and as a genetic guide in the selection of the parents for hybridization.

## REFERENCES

- ACCIARRI N., RESTAINO F., VITELLI G., PERRONE D., ZOTTINI M., PANDOLFINI T., SPENA A., ROTINO G.L., 2002. Genetically modified parthenocarpic eggplants: improved fruit productivity under both greenhouse and open field cultivation. *Biotechnol. 2*: <http://www.biomedcentral.com/1472-6750/2/4>.

- ALBA V., LOTTI T., D'ALESSANDRO A., MENNELLA G., RICIARDI L., SUNSERI F., 2005. Genetic diversity on African eggplant: molecular and chemical analyses. Proc. XLIX Italian Soc. Agric. Gen. Ann. Congress, 12-15 Sept., Potenza, Italy: 135-139.
- ANASO H.U., 1991. Comparative cytological study of *Solanum aethiopicum* Gilo group, *Solanum aethiopicum* Shum group and *Solanum anguivi*. Euphytica 53(2): 81-85.
- ANASO H.U., OKEREKE G.U., UZO J.O., 1990. Breeding for high yield of berries and improved yield attributes in *Solanum aethiopicum* L. Tropical Agric. 67(4): 371-5.
- ANO G., 1989. The interspecific hybridization, a source of variability: an example the eggplant. Acta Hort. 242: 151-154.
- BUKENYA Z.R., 1994. *Solanum macrocarpon*: an underutilised but potential vegetable in Uganda. Proceedings XIIIth Plenary Meeting AETFAT (J.H. Senyani and A.C. Chikuni, eds), 17-24, Malawi.
- BUKENYA Z.R., CARASCO J.F., 1994. Biosystematic study of *Solanum macrocarpon* – *S. dasyphyllum* complex in Uganda and relations with *Solanum linnaeanum*. East African Agric. For. J. 59(3): 187-204.
- BUKENYA Z.R., HALL J.B., 1987. Six cultivars of *Solanum macrocarpon* (*Solanaceae*) in Ghana. Euphytica 17(1): 91-95.
- CARUSO G., 2001. Morpho-physiological characterization of *Solanum aethiopicum* L. Gilo group. Agric. Mediterr. 131(3-4): 153-163.
- CARVALHO A.C.P.P., RIBEIRO R.L.D., 2002. Analise da capacidade combinatoria em cruzamentos dialelicos de tres cultivares de jiloeiro. Hort. Brasil. 20(1): 48-51.
- CHOUDHURY B., 1976. Vegetables. 4<sup>th</sup> Revised Edition National Book Trust, New Delhi, India.
- COBORU, 2006. Research Centre for Cultivar Testing. [www.coboru.pl](http://www.coboru.pl)
- COLLONNIER C., FOCK I., KASHYAP V., ROTINO G.L., DAUNAY M.C., LIAN Y., MARISKA I.K., RAJAM M.V., SERVAES A., DUCREUX G., SIHACHAKR D., 2001. Applications of biotechnology in eggplant. Plant Cell, Tiss. Org. Cult. 65(2): 91-107.
- DAUNAY M.C., CHAPUT M.H., SIHACHAKR D., ALLOT M., VEDEL F., DUCREUX G., 1993. Production and characterization of fertile somatic hybrids of eggplant (*Solanum melongena* L.) with *Solanum aethiopicum* L. Theor. Appl. Genet. 85, 6-7: 841-850.
- DAUNAY M.C., GEBHARDT C., HENNART J.W., JAHN M., LESTER R.N., 2006. Genetic resources of eggplant (*Solanum melongena* L.) and allied species: a new challenge for molecular geneticists and eggplant breeders. Plant and Animal Genome XIV Conference, January 14-18, 145-139, San Diego, California.

- DAUNAY M.C., LESTER R.N., 1988. The usefulness of taxonomy for *Solanaceae* breeders, with special reference to the genus *Solanum* and to *Solanum melongena* L. (eggplant). *Cap. New.* 7: 70-79.
- DAUNAY M.C., MAGGIONI L., LIPMAN E., 2003. European Cooperative Programme For Crop Genetic Resources Network, *Solanaceae* Genetic Resources in Europe. Report of two meetings: 21 September 2001, Nijmegen, The Netherlands and 22 May 2003, Skierniewice, Poland.
- DOGANLAR S., FRARY A., DAUNAY M.C., LESTER R.N., TANKSLEY S.D., 2002a. A comparative genetic linkage map of eggplant (*Solanum melongena*) and its implications for genome evolution in the *Solanaceae*. *Genetics* 161: 1697-1711.
- DOGANLAR S., FRARY A., DAUNAY M.C., LESTER R.N., TANKSLEY S.D., 2002b. Conservation of gene function in the *Solanaceae* as revealed by comparative mapping of domestication traits in eggplant. *Genetics* 161: 1712-1726.
- DONZELLA G., SPENA A., ROTINO G.L., 2000. Transgenic parthenocarpic eggplants: superior germplasm for increased winter production. *Mol. Breed.* 6(1): 79-86.
- ECP/GR EGGPLANT DATABASE, 2006. <http://www.bgard.science.ru.nl/WWW-IPGRI/eggplant.htm>.
- FAOSTAT Data, 2006. Agricultural Data. <http://faostat.fao.org>.
- FAROOQUI M.A., RAO A.V., JAYASREE T., SADANANDAM A., 1997. Induction of atrazine resistance and somatic embryogenesis in *Solanum melongena*. *Theor. Appl. Genet.* 95: 702-705.
- FRANKLIN G., LAKSHMI SITA G., 2003. *Agrobacterium tumefaciens*-mediated transformation of eggplant (*Solanum melongena* L.) using root explants. *Plant Cell Rep.* 21: 549-554.
- FRARY A., DOGANLAR S., DAUNAY M.C., TANKSLEY S.D., 2003. QTL analysis of morphological traits in eggplant and implications for conservation of gene function during evolution of solanaceous species. *Theor. Appl. Genet.* 107: 359-370.
- FURINI A., WUNDER J., 2003. Analysis of eggplant (*Solanum melongena*)-related germplasm: morphological and AFLP data contribute to phylogenetic interpretations and germplasm utilization. *Theor. Appl. Gen.* 108(2): 197-208.
- GOWDA P.H.R., SHIVASHANKAR K.T., JOSHI S.H., 1990. Interspecific hybridization between *Solanum melongena* and *Solanum macrocarpon*: study of the F<sub>1</sub> hybrid plants. *Euphytica* 48(1): 59-61.

- JARL C.I., RIETVELD E.M., DE HAAS J.M., 1999. Transfer of fungal tolerance through interspecific somatic hybridisation between *Solanum melongena* and *S. torvum*. *Plant Cell Rep.* 18: 791-796.
- KARIHALOO J.L., BRAUNER S., GOTTLIEB L.D., 1995. Random amplified polymorphic DNA variation in the eggplant, *Solanum melongena* L. (*Solanaceae*). *Theor. Appl. Genet.* 90(6): 767-770.
- KARIHALOO J.L., GOTTLIEB L.D., 1995. Allozyme variation in the eggplant, *Solanum melongena* L. (*Solanaceae*). *Theor. Appl. Genet.* 90(3-4): 578-583.
- KARIHALOO J.L., KAUR M., SINGH S., 2002. Seed protein diversity in *Solanum melongena* L. and its wild and weedy relatives. *Genet. Res. Crop Evolution* 49(9): 533-539.
- KASHYAP V., VINOD KUMAR S., COLLONNIER C., FUSARI F., HAICOUR R., ROTINO G.L., SIHACHAKR D., RAJAM M.V., 2003. Biotechnology of eggplant. *Scientia Hort.* 97(1): 1-25.
- KUMAR P.A., MANDAOKAR A., SREENIVASU K., CHAKRABARTI S.K., BISARIA S., SHAEMA S.R., KAUR S., SHARMA R.P., 1998. Insect-resistant transgenic brinjal plants. *Mol. Breed.* 4(1): 33-37.
- LESTER R.N., 1986. Taxonomy of scarlet eggplants, *Solanum aethiopicum* L. *Acta Hort.* (ISHS) 182: 125-132.
- LESTER R.N., 1998. Genetic resources of capsicum and eggplants. Xth EUCARPIA Meeting on Genetic and Breeding of Capsicum and Eggplant. 25-30, Avignon, France.
- LESTER R.N., DAUNAY M.C., 2003. Diversity of African vegetable *Solanum* species and its implications for a better understanding of plant domestication. Knüpffer, H. and Ochsmann, J. (eds); *Rudolf Mansfeld and plant genetic resources; Schriften zu Genetischen Ressourcen (DEU) Symposium dedicated to the 100th birthday of Rudolf Mansfeld; Gatersleben (DEU)*, 22: 137-152.
- LEVIN R.A., MYERS N.R., BOHS L., 2006. Phylogenetic relationship among the "spiny solanums" (*Solanum* subgenus *Leptostemonum*, *Solanaceae*). *Am. J. Bot.* 93: 157-169.
- LEVIN R.A., WATSON K., BOHS L., 2005. A four-gene study of evolutionary relationships in *Solanum* section *Acanthophora*. *Am. J. Bot.* 92(4): 603-612.
- MACE E.S., LESTER R.N., GEBHARDT C.G., 1999. AFLP analysis of genetic relationships among the cultivated eggplant, *Solanum melongena* L., and wild relatives (*Solanaceae*). *Theor. Appl. Genet.* 99: 626-633.
- MACHA E.S., 2005. African eggplants promising vegetables for home consumption and sale in Tanzania. *Proceedings of the Third Horticulture Workshop on Sustainable Horticultural Production in the Tropics*, 26-29 Nov. 2003, Maseno, Kenya.

- MAGIOLI C., MANSUR E., 2005. Eggplant (*Solanum melongena* L.): tissue culture, genetic transformation and use as an alternative model plant. *Acta Bot. Bras.* 19(1): 139-148.
- MAGIOLI C., ROCHA A.P.M., OLIVEIRA D.E., MANSUR E., 1998. Efficient shoot organogenesis of eggplant (*Solanum melongena* L.) induced by thidiazuron. *Plant Cell Rep.* 17: 661-663.
- MAGIOLI C., ROCHA A.P.M., TARRE E., SANTIAGO-FERNANDES L.D., OLIVEIRA D.E., KRUL W.R., MASUR E., 2001. Effect of morphological factors, antibiotics and *Agrobacterium* co-cultivation in the efficiency of somatic embryogenesis of eggplant (*Solanum melongena* L.). *J. Plant Biotechnol.* 3(1): 19-25.
- MARTIN F.W., RHODES A.M., 1979. Subspecific grouping of eggplant cultivars. *Euphytica* 28: 367-383.
- MUELLER L.A., SOLOW T.H., TAYLOR N., SKWARECKI B., BUELS R., BINNS J., LIN CH., WRIGHT M., AHRENS R., WANG Y., HERBST E.V., KEYDER E.R., MENDA N., ZAMIR D., TANKSLEY D., 2005. The SOL Genomic Network. A comparative resource for *Solanaceae* biology and beyond. *Plant Physiol.* 138: 1310-1317.
- OLMSTEAD R.G., PALMER J.D., 1997. Implications for the phylogeny, classification, and biogeography of *Solanum* from cpDNA restriction site variation. *Systematic Bot.* 22(1): 19-29.
- OMIDIJI M.O., 1983. Evaluation of some F<sub>1</sub> hybrids and cultivars of the local eggplant, *Solanum gilo* Raddi in South-Western Nigeria. *Acta Hort.* 123: 91-98.
- PESSARAKLI M.M., DRIS R., 2004. Pollination and breeding of eggplants. *Food, Agric. Environ.* 2(1): 218-219.
- PRABHAVATHI V., YADAV J.S., KUMAR P.A., RAJAM M.V., 2002. Abiotic stress in transgenic eggplant (*Solanum melongena* L.) by introduction of bacterial mannitol phosphodehydrogenase gene. *Mol. Breed.* 9(2): 137-147.
- PROHENS J., BLANCA J.M., NUEZ F., 2005. Morphological and molecular variation in a collection of eggplants from secondary center of diversity: implications for conservation and breeding. *J. Am. Soc. Hort. Sci.* 130(1): 54-63.
- RAHMAN M.A., RASHID M.A., SALAM M.A., MASUD M.A.T., MASUM A.S.M.H., HOSSAIN M.M., 2002. Performance of some grafted eggplant genotypes on wild *Solanum* root stocks against root-knot nematode. *OnLine J. Biol. Sci.* 2(7): 446-448.
- RIZZA F., MENNELLA G., COLLONNIER C., SIHACHAKR D., KASHYAP V., RAJAM M., PRESTERA M., ROTINO G., 2002. Androgenic dihaploids from somatic hybrids between *Solanum melongena* and *S. aethiopicum* group gilo as a source of resistance to *Fusarium oxysporum* f. sp. *melongenae*. *Plant Cell Rep.* 20(11): 1022-1032.



- SAKATA Y., LESTER R.N., 1997. Chloroplast DNA diversity in brinjal eggplant (*Solanum melongena* L.) and related species. *Euphytica* 97: 295-301.
- SCHAFF D.A., JELENKOVIC G., BOYER C.D., POLLACK B.L., 1982. Hybridization and fertility of hybrid derivatives of *Solanum melongena* L. and *Solanum macrocarpon* L. *Theor. Appl. Genet.* 62(2): 149-153.
- SEEDQUEST, 2006. Information Services for Seeds Professional. [www.seedquest.com](http://www.seedquest.com).
- SEEK A., 1997. Conservation and utilization of germplasm of traditional African vegetables in Senegal. In: *Traditional African vegetables*. Guarino L. (ed.). Proceedings of the IPGRI International Workshop on Genetic Resources of Traditional Vegetables in Africa: Conservation and Use, 29-31 August, Nairobi, Kenya.
- SHARMA P., RAJAM M.V., 1995. Genotype, explants and position effects on organogenesis and embryogenesis in eggplant (*Solanum melongena* L.). *J. Exp. Bot.* 46: 13-141.
- SIDHU A.S., BAL S.S., BEHERA T.K., RANI M., 2005. An outlook in hybrid eggplant breeding. *J. New Seeds* 6 (2/3): 15-29.
- SINGH A.K., SINGH M., SINGH A.K., SINGH R., KUMAR S., KALLOO G., 2006. Genetic diversity within the genus *Solanum* (*Solanaceae*) as revealed by RAPD markers. *Current Sci.* 90(5): 711-716.
- SOL GENOMIC NETWORK, 2006. <http://sgn.cornell.edu>.
- STEDJE B., BUKENYA-ZIRABA R., 2003. RAPD variation in *Solanum anguivi* Lam. and *S. aethiopicum* L. (*Solanaceae*) in Uganda. *Euphytica* 131(3): 293-297.
- SWARUP V., 1995. Genetic resources and breeding of aubergine (*Solanum melongena* L.). *Acta Hort.* 412: 71-79.

## OBERŻYNA – POCHODZENIE, KIERUNKI HODOWLI I ZASOBY GENETYCZNE, PRACA PRZEGLĄDOWA

Streszczenie: Oberżyna (*Solanum melongena* L.) jest ważnym gospodarczo gatunkiem z rodziny *Solanaceae*, pochodzącym z południowych Indii, a szeroko uprawianym w Ameryce Północnej i Południowej, Europie i Azji. Spokrewnione z tym gatunkiem *S. aethiopicum* i *S. macrocarpon* pochodzą z Afryki i tam są uprawiane. Oberżyna jest nowym warzywem w Polsce, do niedawna uprawianym tylko w warunkach szklarniowych. Hodowla nowych odmian i doskonalenie technologii produkcji spowodowały zwiększenie areалу upraw i wzrost popularności oberżyny wśród konsumentów. W publikacji przedstawiono szczegółową systematykę oberżyny i gatunków pokrewnych, ich pochodzenie oraz historię uprawy. Zestawiono informacje na temat zasobów genetycznych na świecie i najważniejszych kierunków hodowli gatunku. Zaprezentowano wybrane grupy odmian i charakterystykę najważniejszych kreacji hodowlanych.

Received November 27, 2006; accepted May 5, 2007

Cultivated eggplants " origin, breeding objectives and genetic resources, a review. *Folia Horticulturae* Ann. 19/1 :97-114. Genetic Variability in Eighteen Cultivars of *Solanum anguivi* Lam. using Principal Component Analysis (PCA) and Single Linkage Cluster Analysis (SLCA). *Annals of Biological Research*, 2 (4): 62-67. [10]. Nowadays eggplant is cultivated in tropical and subtropical regions of the world; China is the top producer while the Philippines is in the 9th place. Jorge R. Sahagun. References: Daunay, M.-C. and Chadha, M.L. 2004. *Solanum melongena* L. [Internet] Record from Protabase. Grubben, G.J.H. and Denton, O.A. (Editors). PROTA (Plant Resources of Tropical Africa / Ressources vgtales de l'Afrique tropicale), Wageningen, Netherlands. < <http://database.prota.org/search.htm>>. Accessed 13 September 2011. Cultivated eggplants origin, breeding objectives and genetic resources, a review. *FOLIA HORTICULTURAE* Ann. 19/1, 2007, 97-114 Wang J.-X., Gao T.-G., and Knapp S. 2008. Therefore, genetics of fruit phenolics, fruit flesh colour, and browning related traits were studied in the genotypes of eggplant, comprising of nine cultivated varieties and one accession of eggplant's primary genepool wild relative *Solanum insanum* (INS2). These accessions were genotyped based on the 7335 polymorphic single-nucleotide polymorphisms (SNP) markers. After that, genotypes were crossed in half diallel fashion to produce 45 hybrids. The INS2 displayed the highest values for the total phenolics and chlorogenic acid content (CGA). Sekara, A.; Cebula, S.; Kunicki, E. Cultivated eggplants "Origin, breeding objectives and genetic resources, a review. *Folia Hortic.* 2007, 19, 97"114. Genetics, Genomics and Breeding of Peppers and Eggplants. p. 116. CrossRef. Google Scholar. Dias, Jo"o Silva and Ortiz, Rodomiro 2015. *Sustainable Agriculture Reviews*. Vol. 17, Issue. , p. 121. CrossRef. Google Scholar. Ying, Yangwei Jin, Wei Yu, Bingwen Lv, Shaowu Wu, Xiaofei Yu, Haixiang Shan, Jin Zhu, Dan Jin, Qinhan and Mu, Ying 2016. Support vector machine classification for determination of geographical origin of Chinese ginseng using microwave plasma torch-atomic emission spectrometry. *Analytical Methods*, Vol. 8, Issue. 25, p. 5079. S"m"ara A., Cebula S., Kunicki E. (2007): Cultivated eggplants " origin, breeding objectives and genetic resources: A review. *Folia Horticulturae*, 19: 97"114. S"m"ara A., Kalisz A., B"czek-Kwinta R., Gaw"da M., Pohl A., Grabowska A. (2015): The effect of abiotic stresses applied in the juvenile phase of eggplant ontogeny on chemical composition of seedlings and fruits. Swarup V. (1995): GENETIC RESOURCES AND BREEDING OF AUBERGINE (*SOLANUM MELONGENA* L.). *Acta Horticulturae*, , 71-79 <https://doi.org/10.17660/ActaHortic.1995.412.6>. Syrenius S. (1613): Zielnik. Krakow, Printing House of Basilii Skalski.